

BIO-BASED SUCCINIC ACID FOR EXFOLIATION

CREATING VALUE WITH NATURAL PERSONAL CARE INGREDIENTS THAT PERFORM

Bio-based succinic acid can be used as an effective exfoliating agent. BioAmber's bio-based succinic acid is naturally derived from non-GMO plant sugar, and does not fall into the alpha-hydroxy acid category. It has been shown to be equally effective in exfoliating dry skin compared to glycolic acid and lactic acid.



BIO-SA™ NATURAL PERSONAL CARE GRADE INCI NAME: BUTANEDIOIC ACID

ANALYSIS	UNIT	SPECIFICATIONS
Molecular weight	g/mol	118.09
CAS #	---	110-15-6
Appearance	---	White crystalline powder, Odorless, characteristics acid taste
Particle size	---	150µm -800µm
Melting Point	---	186°C - 190°C
Water content	%	≤ 0.5%
Assay (USP Monograph)	%	≥ 99.0%
Residue on ignitions (USP Method 281)	%	< 0.025%
Other organic acids	%	< 0.5%
Lead (EPA Method 60108) * Formaldehyde free	µg/g	< 2

Potential applications

- Exfoliation products (cell turnover)
- Skin peeling products
- Creams, serums, lotions, cleansers and toners
- Moisturizing skin creams and lotions

Irritation / Sensitization testing

- The rinse-off succinic acid gel was tested for skin irritation in a 21 day cumulative irritation study with 20 subjects.
- The rinse-off gels were diluted to 20% in water and applied five days weekly for 21 days to the same site.
- No irritation response was observed for any subject with the rinse-off succinic acid gels. (Tests performed by AMA Laboratories)




NORTH AMERICA :: +1.514.844.8000
info@bio-amber.com

™ Trademark owned by BioAmber Inc. © 2015 BioAmber Inc. All Rights Reserved.

www.bio-amber.com

BIO-BASED SUCCINIC ACID FOR EXFOLIATION

CREATING VALUE WITH NATURAL PERSONAL CARE INGREDIENTS THAT PERFORM



EXFOLIATION GEL FORMULATION

(Prepared by Cosmetech Laboratories)

	INGREDIENT (INCI)	WEIGHT PERCENT
PHASE A	Water	58.95
	Guar Gum	0.50
	Xanthan Gum	0.40
PHASE B	Potassium Sorbate	0.10
	Phenoxyethanol	0.70
PHASE C	Water	25.00
	Succinic Acid	7.50
PHASE C1	50% NaOH in water	6.60
PHASE D	Sodium Chloride	0.25

MANUFACTURING INSTRUCTIONS

Phase A: Disperse gums in water one at a time and mix until they are fully hydrated.


Phase B: Add Phase B to Phase A.

Phase C: Premix Phase C in a separate vessel. Use Phase C1 to adjust pH of phase C to 4.0-4.50 and mix until the succinic acid dissolves. Warm up to 45-50 °C if necessary. Add phase C/C1 to batch and mix until uniform.

Phase D: Add Phase D to batch and mix until uniform.

Efficacy

- Rinse-off exfoliating gels were prepared containing 7.5% by weight of bio-based succinic acid, 10.77% glycolic acid (70%), and 8.33% lactic acid (90%). The rinse-off exfoliating gels were applied to a wet forearm.
- The area was rubbed gently for 30 seconds. The gel was left on the forearm for 30 minutes, then rinsed and patted dry. D-Squame surface testing was conducted prior to treatment, and 15 minutes after rinse-off.
- Gels containing bio-based succinic acid at higher pH decreased the surface scaliness of skin more effectively than the traditional alpha-hydroxy acids at lower pH.



+ Exfoliating efficiency through skin debris collection via D-Squame surface sampling. Average of five female test subjects ages 41-45. Testing for exfoliation was conducted at AMA Laboratories



NORTH AMERICA :: +1.514.844.8000
info@bio-amber.com

™ Trademark owned by BioAmber Inc. © 2015 BioAmber Inc. All Rights Reserved.

www.bio-amber.com